

# Cloud modeling of a quiet solar region in $H\alpha$

Z. F. Bostanci<sup>1,2</sup> and N. Al Erdoğan<sup>1</sup>

<sup>1</sup> Istanbul University, University Observatory, 34119 University-Istanbul, Turkey

<sup>2</sup> Sterrekundig Instituut Utrecht, Postbus 80 000, NL3508 TA Utrecht, The Netherlands  
e-mail: fbostanci@gmail.com

**Abstract.** We present chromospheric cloud modeling on the basis of  $H\alpha$  profile-sampling images taken with the Interferometric Bidimensional Spectrometer (IBIS) at the Dunn Solar Telescope (DST). We choose the required reference background profile by using theoretical NLTE profile synthesis. The resulting cloud parameters are converted into estimates of physical parameters (temperature and various densities). Their mean values compare well with the VAL-C model.

**Key words.** Line: profiles – Techniques: spectroscopic – Sun: chromosphere

## 1. Introduction

The solar chromosphere observed in  $H\alpha$  shows a mass of fibrillar structures. They are called mottles when seen on the disk, spicules when seen at the limb. Studies of these structures are important to understand chromospheric dynamics and its contribution to outer-atmosphere heating.

Chromospheric observations sampling spectral profiles give the opportunity to derive physical properties per fine structure. We do that here for  $H\alpha$  using the DST/IBIS data of Cauzzi et al. (2009). Cloud modeling following Beckers (1964) is the usual method to obtain line formation parameters by matching the observed contrast profile of a structure with a theoretical one (see review by Tziotziou 2007). This approach can only be used if the studied structure is fully separated from the underlying atmosphere, and necessitates description of the latter by a background profile.


Its choice or determination is crucial (see e.g. Durrant 1975). We address this issue by trying different synthetic  $H\alpha$  profiles. We then use the method of Tsiropoulou & Schmieder (1997) to derive physical parameters from the cloud model results.

## 2. Observations

In March 2007 a quiet-sun area near disk center was observed in  $H\alpha$  with IBIS at the DST (Cavallini 2006; Reardon & Cavallini 2008). These observations were presented and analyzed by Cauzzi et al. (2009). The  $H\alpha$  line was sampled at 24 spectral positions at step intervals of 90 mÅ in  $H\alpha$  in a sequence of 192 spectral scans at a cadence of 15.4 seconds. Line profiles were constructed for each pixel in the field of view (diameter 80 arc-sec). For each spectral profile, the line-center wavelength was established by fitting a polynomial to the five spectral samplings with least intensity. The minimum of the fit defines the per-pixel intensity minimum and line-of-sight

---

Send offprint requests to: Z. F. Bostanci




**Fig. 1.** *Upper image:* profile-minimum intensity. *Lower image:* profile-minimum Dopplershift, with blueshift black, redshift white.

velocity. Figure 1 shows the minimum intensity and Doppler velocity images from a single scan taken at one of the best seeing moments. We refer to Cauzzi et al. (2009) for more detail.

### 3. Results & Conclusions

#### 3.1. Cloud Model

The traditional cloud model delivers the four parameters source function  $S$ , optical thickness at line center  $\tau_0$ , Doppler width  $\Delta\lambda_D$ , and line of sight velocity  $v_{LOS}$ . The model assumes an optically thin, homogeneous cloud that is illuminated by uniform radiation from below,



**Fig. 2.** Background profiles, with the intensities normalized to the continuum value. *Solid:* Kurucz model. *Dashed:* FAL-C model. *Dotted:* observed mean profile.

so that these parameters are assumed constant along the line of sight through the cloud. The observed contrast profiles are then matched with theoretical contrast profiles given by:

$$\frac{I(\lambda) - I_0(\lambda)}{I_0(\lambda)} = \left( \frac{S}{I_0(\lambda)} - 1 \right) (1 - e^{-\tau(\lambda)}), \quad (1)$$


where  $I(\lambda)$  is the local profile,  $I_0(\lambda)$  the reference background profile and  $\tau(\lambda)$  the optical thickness

$$\tau(\lambda) = \tau_0 \exp \left[ - \left( \frac{\lambda - \lambda_c(1 - v_{LOS}/c)}{\Delta\lambda_D} \right)^2 \right]. \quad (2)$$

The parameter fitting is achieved by iterative least-square matching of the observed contrast profile with a theoretical one.

#### 3.2. Background profile


The background profile  $I_0(\lambda)$  represents the irradiation from supposedly plane-parallel atmosphere underlying the cloud-like chromospheric structure. Its choice has a significant effect on the resulting cloud parameters. We here present H $\alpha$  cloud modeling results for the IBIS scan with the best seeing using three different background profiles: a synthetic profile computed with a one-dimensional NLTE line formation code from the FAL-C (Fontenla et al.



**Fig. 3.** Occurrence distributions of the cloud parameters. Clockwise source function  $S$ , optical thickness at line center  $\tau_0$ , Doppler width  $\Delta\lambda_D$ , and line of sight velocity  $v_{LOS}$ . *Solid*: Kurucz background profile. *Dashed*: FAL-C background profile. *Dotted*: observed mean profile used as background profile.

1993) standard model which contains a chromospheric temperature rise, a synthetic NLTE profile similarly computed from the Kurucz (Kurucz 1979, 1992a,b) radiative-equilibrium model in which the temperature declines outward without chromosphere, and the spatial-temporal average of all observed  $H\alpha$  profiles over the full field of view during the whole 50-min time series. The three profiles are shown in Figure 2.

With each background profile, cloud-model fitting was applied to the observed  $H\alpha$  profile at all  $1.74 \times 10^5$  pixels in the IBIS field of view. We rejected the pixels with resulting values  $S > 0.8I_c$  (in units of continuum intensity) and also the pixels giving  $\tau_0 > 5$ . In addition, the cloud model routine did not converge or did not yield physically acceptable values for 0.04%, 7.47% and 23.19% of the total when we used the Kurucz, FAL-C, and observed mean profile, respectively. Figure 3 displays the remaining distributions of the cloud model parameters for each background profile. It shows that using the Kurucz profile permits a



**Fig. 4.** Scatter plots. *Left column*:  $H\alpha$  source function  $S$  against the observed profile-minimum intensity, respectively with the FAL-C background profile (top), the Kurucz background profile (middle), and the observed mean profile as background profile (bottom). *Right column*: idem for the fitted line-of-sight cloud velocity  $v_{LOS}$  against the observed profile-minimum Dopplershift. The numbers at the upper-right corners specify the overall Pearson correlation coefficient.

solution for many more  $H\alpha$  profiles, with narrower parameter distributions.

Figure 4 shows scatter plots for the resulting values of  $S$  and  $v_{LOS}$  against the observed profile-minimum intensity and Dopplershift, respectively, when using the three different background profiles. The best correlations are found between these cloud parameters and profile-minimum measurements when the Kurucz synthetic profile is used.

These comparisons suggest that the synthetic Kurucz profile is the best choice as background profile for cloud modeling of these  $H\alpha$  observations. The spread between these tests

**Table 1.** Comparison of the observed parameters with the values inferred from VAL-C atmosphere models.

Physical Parameters	Quiet Chromosphere	
	Observational	VAL-C
$S$ ( $I_c$ )	$0.19 \pm 0.02$	–
$\tau_0$	$1.50 \pm 0.46$	–
$\Delta\lambda_D$ (Å)	$0.46 \pm 0.04$	–
$v_{LOS}$ (km s $^{-1}$ )	$-1.74 \pm 3.37$	–
$N_1$ ( $10^{10}$ cm $^{-3}$ )	$2.21 \pm 0.42$	1.24
$N_2$ ( $10^4$ cm $^{-3}$ )	$2.54 \pm 0.88$	2.88
$N_e$ ( $10^{10}$ cm $^{-3}$ )	$5.13 \pm 0.94$	3.54
$N_H$ ( $10^{10}$ cm $^{-3}$ )	$8.02 \pm 1.47$	4.67
$T$ ( $10^4$ K)	$1.43 \pm 0.40$	1.07
$P$ (dyn cm $^{-2}$ )	$0.27 \pm 0.09$	0.13
$M$ ( $10^{-5}$ gr cm $^{-2}$ )	$3.61 \pm 0.55$	0.62
$\rho$ ( $10^{-13}$ gr cm $^{-3}$ )	$1.75 \pm 0.28$	1.09
$\chi_H$	$0.63 \pm 0.01$	–

confirms that the issue of the selection of a background profile is a key one in chromospheric cloud modeling.

### 3.3. Physical parameters

We then applied cloud model fitting using the Kurucz background profile to all 192 spectral scans in the 50-min time series. Afterwards, we converted the resulting cloud parameters into more physical parameters with the method of Tsiropoula & Schmieder (1997). The resulting mean values and standard deviations are given in Table 1. For comparison the values of the VAL-C atmosphere model of (Vernazza et al. 1981) at the height where its  $N_2$  population is close to the mean value in the cloud determinations are also listed. Table 1 shows good agreement between the cloud model and VAL-C values.

*Acknowledgements.* We are indebted to Kevin Reardon for supplying the observations and explanation, to Han Uitenbroek for suggesting and computing the theoretical background profiles, to Rob Rutten for improvement of this paper, and to all three and Gianna Cauzzi and Alexandra Tritschler for discussions. This research project has been supported by a USO–SP Marie Curie Early Stage Research Training Fellowship from the European Community under contract num-

ber MEST-CT-2005-020395. IBIS was built by INAF/Osservatorio Astrofisico di Arcetri with contributions from the Università di Firenze and the Università di Roma Tor Vergata. IBIS construction and operation has been supported by the Italian Ministero dell’Università e della Ricerca (MUR), as well as the Italian Ministry of Foreign Affairs (MAE). NSO is operated by the Association of Universities for Research in Astronomy, Inc. (AURA), under cooperative agreement with the National Science Foundation. This work was supported by the Research Fund of Istanbul University as project number 851.

### References

- Beckers, J. M. 1964, A Study of the Fine Structures in the Solar Chromosphere (AFCRL Environmental Research Paper No. 49, PhD thesis Utrecht University)
- Cauzzi, G., Reardon, R., Rutten, R. J., Tritschler, A., Uitenbroek, H. 2009, A&A, 503, 577
- Cavallini, F. 2006, Sol. Phys., 236, 415
- Durrant, C.J. 1975, Sol. Phys., 44, 41
- Fontenla, J. M., Avrett, E. H., & Loeser, R. 1993, ApJ, 406, 319
- Kurucz, R. L. 1979, ApJS, 40, 1
- Kurucz, R. L. 1992a, Rev. Mex. Astron. Astrofis., 23, 181
- Kurucz, R. L. 1992b, Rev. Mex. Astron. Astrofis., 23, 187
- Reardon, K. P., Cavallini, F. 2008, A&A, 481, 897
- Tsiropoula, G., Schmieder, B. 1997, A&A, 324, 1183
- Tziotziou, K. 2007, in The Physics of Chromospheric Plasmas, ed. P. Heinzel, I. Dorotovič, & R. J. Rutten, PASPC, 368, 217
- Vernazza, J. E., Avrett, E. H., Loeser, R. 1981, ApJS, 45, 635